

ON ATKIN-LEHNER QUOTIENTS OF SHIMURA CURVES

BRUCE W. JORDAN AND RON LIVNÉ

ABSTRACT. We study the Čerednik-Drinfeld p -adic uniformization of certain Atkin-Lehner quotients of Shimura curves over $\mathbf{Q}$. We use it to determine over which local fields they have rational points and divisors of a given degree. Using a criterion of Poonen and Stoll we show that the Shafarevich-Tate group of their jacobians is not of square order for infinitely many cases.

In [PSt] Poonen and Stoll have shown that if the Shafarevich-Tate group of a principally polarized abelian variety A over a global field K is finite, then its order can be twice a square as well as a square. They call A/K even if the quotient of its Shafarevich-Tate group by the maximal divisible subgroup has order a square and odd otherwise. They prove [PSt, Corollary 10] that the jacobian of a curve X/K of genus g is odd if and only if the number of places v of K where X fails to have a K_v -rational divisor of degree $g-1$ is odd. They also show that infinitely many hyperelliptic jacobians over $\mathbf{Q}$ are odd for every even genus.

Let B be an indefinite rational quaternion division algebra of discriminant $\text{Disc } B$. Let $V_B/\mathbf{Q}$ be the Shimura curve corresponding to a maximal order in B . In [JL1] and [JL2] we determined when Shimura curves V_B have rational points, rational divisors of a given degree, and rational divisor classes of a given degree over any local field. The key ingredient of our analysis was the explicit p -adic uniformization of these curves given by Drinfeld [Dr]. The criterion of Poonen and Stoll then immediately yields that the jacobian of $V_B/\mathbf{Q}$ is even ([PSt, Theorem 23]). Atkin-Lehner involutions w_d for $d \mid \text{Disc } B$ act on the Shimura curves V_B . Set $V_B^{(d)} = V_B^{(d)}/\mathbf{Q} = V_B/w_d$. We will show that these Atkin-Lehner quotients $V_B^{(d)}/\mathbf{Q}$, in contrast to $V_B/\mathbf{Q}$, yield examples of odd nonhyperelliptic jacobians of arbitrarily high genus. For simplicity we consider quaternion algebras B in which exactly two odd primes ramify and study the Atkin-Lehner quotient $V_B^{(p)}/\mathbf{Q}$ where p is a prime. For a related study of more general Shimura curves over totally real fields see [JLV]. In this paper we will prove the following:

Date: February 4, 1998.

1991 Mathematics Subject Classification. 11G18, 11G20, 14G20, 14G35, 14H40.

Key words and phrases. Shimura curves, Atkin-Lehner involutions, Shafarevich-Tate groups, p -adic uniformization.

The first author was partially supported by grants from the NSF and PSC-CUNY.

Both authors were supported by a joint Binational Israel-USA Foundation grant.

Theorem 1. *Let $p \neq q$ be primes with $p \equiv 5 \pmod{24}$ and $q \equiv 5 \pmod{12}$. Furthermore suppose that p is not a square modulo q . Let B be the indefinite rational quaternion algebra with $\text{Disc } B = pq$. Then the jacobian of $V_B^{(p)}/\mathbf{Q}$ is odd.*

The genus of the curves $V_B^{(p)}$ in Theorem 1 goes to infinity with $\text{Disc } B$ and we show at the end of Section 2 that only finitely many of these curves can be hyperelliptic.

1. LOCAL DIOPHANTINE PROPERTIES OF ATKIN-LEHNER QUOTIENTS

Let B be an indefinite rational quaternion division algebra of discriminant pq , where p and q are distinct odd primes. Put $V^{(p)}/\mathbf{Q} = V_B/w_p$ and $V^{(q)}/\mathbf{Q} = V_B/w_q$. For a curve $X/\mathbf{Q}$, an extension field $K/\mathbf{Q}$, and an integer r we denote by $\text{Pic}^r X(K)^+$ the set of divisor classes on X of degree r containing a K -rational divisor. Let $B^{p\text{-int}}/\mathbf{Q}$ be the definite quaternion algebra obtained from B by interchanging local invariants at p and ∞ ; the reduced discriminant of $B^{p\text{-int}}$ is q . For $\alpha, \beta \in \mathbf{Q}$ denote by $B(\alpha, \beta)$ the rational quaternion algebra with basis $1, i, j$, and k satisfying $i^2 = \alpha, j^2 = \beta$, and $ij = -ji = k$. It is ramified at a prime $p \leq \infty$ if and only if the Hilbert symbol $(\alpha, \beta)_p = -1$.

Theorem 2. 1. $\text{Pic}^1 V^{(p)}(\mathbf{R})^+ \neq \emptyset$ if and only if $\mathbf{Q}(\sqrt{p})$ splits B .

2. $\text{Pic}^1 V^{(p)}(\mathbf{Q}_p)^+ \neq \emptyset$.

3. $\text{Pic}^1 V^{(q)}(\mathbf{Q}_p)^+ \neq \emptyset$ if and only if

$$(1) \quad B^{p\text{-int}} \cong B(-1, -pq) \quad \text{or} \quad B^{p\text{-int}} \cong B(-p, -q) .$$

Proof: By applying Shimura's explicit law for the action of complex conjugation on $V_B(\mathbf{C})$, Ogg proved in [Ogg] that $V^{(p)}(\mathbf{R}) \neq \emptyset$ if and only if $\mathbf{Q}(\sqrt{p})$ splits B . It is obvious that $V^{(p)}(\mathbf{R}) \neq \emptyset$ if and only if $\text{Pic}^1 V^{(p)}(\mathbf{R})^+ \neq \emptyset$, proving (1).

For (2) fix a maximal $\mathbf{Z}[1/p]$ -order $\mathcal{D} \subseteq B^{p\text{-int}}$, so that $\mathbf{Z}[1/p]^\times \subseteq \mathcal{D}^\times$. Let N be the reduced norm of $B^{p\text{-int}}$ and let $\text{val}_p(N(\gamma))$ be the p -adic valuation. Define

$$\tilde{\Gamma}_+ = \{\gamma \in \mathcal{D}^\times \mid \text{val}_p(N(\gamma)) \text{ is even}\} \quad \text{and} \quad \Gamma_+ = \tilde{\Gamma}_+/\mathbf{Z}[1/p]^\times .$$

Fix an identification of $B^{p\text{-int}} \otimes \mathbf{Q}_p$ with $\text{Mat}_{2 \times 2}(\mathbf{Q}_p)$. Then $\tilde{\Gamma}_+$ is a subgroup of $\text{GL}_2(\mathbf{Q}_p)$ containing $p = p\text{Id}_{2 \times 2}$ and Γ_+ is a discrete cocompact subgroup of $\text{PGL}_2(\mathbf{Q}_p)$.

Let $M_{\Gamma_+}/\mathbf{Z}_p$ denote the Mumford curve uniformized by $\Gamma_+ \subseteq \text{PGL}_2(\mathbf{Q}_p)$. Drinfeld proves that the curve $V_B \times_{\mathbf{Q}} \mathbf{Q}_p$ is the generic fiber of the w_p -Frobenius quadratic twist of $M_{\Gamma_+}/\mathbf{Z}_p$, see [JL1, Sect. 4]. Hence the quotient $V^{(p)}$ is Mumford uniformized (without a twist) and it has a model over $\mathbf{Z}_p$ whose reduction consists of $\mathbf{F}_p$ -rational $\mathbb{P}^1$'s crossing at $\mathbf{F}_p$ -rational points. In particular over any extension $\mathbf{F}_{p^r}$, $r \geq 2$, there are smooth $\mathbf{F}_{p^r}$ -rational points. By Hensel's lemma (see, e.g., [JL1, Lemma 1.1]) there are points rational over the Witt vectors $W(\mathbf{F}_{p^r})$, $r \geq 2$. The trace to $\mathbf{Q}_p$ of such a point is in $\text{Pic}^r V^{(p)}(\mathbf{Q}_p)^+$. A linear combination using relatively prime r 's (say $r = 2, 3$) gives then a $\mathbf{Q}_p$ -rational divisor of degree 1, proving (2).

For (3), we have $\text{Pic}^2 V_B(\mathbf{Q}_p)^+ \neq \emptyset$ by [JL2, Theorem 2a], so also $\text{Pic}^2 V^{(q)}(\mathbf{Q}_p)^+ \neq \emptyset$. Hence $\text{Pic}^1 V^{(q)}(\mathbf{Q}_p)^+ \neq \emptyset$ if and only if there is a $\mathbf{Q}_p$ -rational effective divisor of odd degree on $V^{(q)}$. Such a divisor is the sum of traces to $\mathbf{Q}_p$ of points in extension fields of

$\mathbf{Q}_p$ at least one of which must be of odd degree over $\mathbf{Q}_p$. Hence we need to show that the condition (1) holds if and only if $V^{(q)}(K) \neq \emptyset$ for some extension $K/\mathbf{Q}_p$ of odd degree. (In fact our proof will show that condition (1) is equivalent to $V^{(q)}(\mathbf{Q}_p) \neq \emptyset$.)

Let Δ be the tree of $\mathrm{SL}(2, \mathbf{Q}_p)$; it is a homogeneous tree of regularity $p+1$. The dual graph (see [Kur], [JL1]) of the special fiber of M_{Γ_+} is given by $G = \Gamma_+ \setminus \Delta$. Let γ_q be an element of the normalizer of $\tilde{\Gamma}_+$ in $\mathcal{D}^\times$ of reduced norm q . Set

$$\tilde{\Gamma}_{+,q} = \langle \tilde{\Gamma}_+, \gamma_q \rangle \subseteq \mathrm{GL}(2, \mathbf{Q}_p) \quad \text{and} \quad \Gamma_{+,q} = \tilde{\Gamma}_{+,q} / \mathbf{Z}[1/p]^\times.$$

From [Dr] it follows that γ_q induces the action of w_q on M_{Γ_+} . Taking the quotients of M_{Γ_+} and V_B by w_q we see that $V^{(q)} \times_{\mathbf{Q}} \mathbf{Q}_p$ is the w_p -Frobenius quadratic twist of the Mumford curve $M_{\Gamma_{+,q}}$ uniformized by $\Gamma_{+,q} \subseteq \mathrm{PGL}(2, \mathbf{Q}_p)$. This gives a model of $V^{(q)} \times_{\mathbf{Q}} \mathbf{Q}_p$ over $\mathbf{Z}_p$ with dual graph $G^{(q)} = w_q \backslash G$ and Frobenius acting as w_p . Let $K/\mathbf{Q}_p$ be an extension with ring of integers $\mathcal{O}_K$ and residue field $\mathbf{k}$. Put $e = e(K/\mathbf{Q}_p)$ and $f = f(K/\mathbf{Q}_p)$. By base change, $V^{(q)} \times_{\mathbf{Q}_p} K$ has an $\mathcal{O}_K$ -model having dual graph $G^{(q)}$ and Frobenius action w_p^f . Moreover, the lengths of edges are multiplied by e , see [JL1, Proposition 3.4].

The graph G is bipartite with w_p interchanging the even and the odd vertices; see, e.g., the discussion preceding Proposition 4-4 in [Kur]. This property is inherited by the quotient graph $G^{(q)}$. We now suppose that $K/\mathbf{Q}_p$ is an extension of odd degree. Then Frobenius acts on $G^{(q)}$ by w_p , which can fix no vertex as it interchanges the even and odd vertices of the bipartition. Hence no component of the special fiber is rational over $\mathbf{k}$, and a point x of $V^{(q)}(K)$ must specialize to a singularity of the special fiber. The length $\ell(r)$ of the edge r of $G^{(q)}$ corresponding to x determines the nature of the singularity at x , and is the order of the stabilizer in $\Gamma_{+,q}$ of any preimage $\tilde{s}$ of it in Δ . As in [JL1, Theorem 5.2], $V^{(q)}(K) \neq \emptyset$ if and only if there is an oriented edge r of $G^{(q)}$ such that $\ell(r)$ is even and $w_p(r)$ is the opposite edge $\bar{r}$. Let s be an edge of G above such an r . Then we see that

- 1) Either $\ell(s)$ is even or $w_q(s) = s$.
- 2) Either $w_p(s) = \bar{s}$ or $w_{pq}(s) = \bar{s}$.

This gives us four cases. The case $\ell(s)$ is even and $w_p(s) = \bar{s}$ is impossible by [JL1, Theorem 5.6] since p and q are both odd. Furthermore if $w_q(s) = s$ and $w_{pq}(s) = \bar{s}$, then we also have $w_p(s) = \bar{s}$. Hence we are reduced to considering two cases. Let $\tilde{s}$ be an edge of Δ lying above s .

We first consider the case $\ell(s)$ is even and $w_{pq}(s) = \bar{s}$. Let γ_{pq} be an element of $\mathcal{D}$ representing w_{pq} . Modifying γ_{pq} by an element of $\mathcal{D}^\times$ we may assume that $N(\gamma_{pq}) = pq$ and $\gamma_{pq}\tilde{s} = \bar{\tilde{s}}$. As in the proof of [JL1, Theorem 5.6] it follows that $B^{p\text{-int}} \cong B(-1, -pq)$. The same argument read in reverse shows that if $B^{p\text{-int}} \cong B(-1, -pq)$ we can get such an edge $\tilde{s}$ which then descends to an edge s of G satisfying $\ell(s) = 2$ and $w_{pq}(s) = \bar{s}$. The edge s of G in turn descends to an edge r of $G^{(q)}$ such that $\ell(r)$ is even and $w_p(r) = \bar{r}$.

Now suppose $w_q(s) = s$ and $w_p(s) = \bar{s}$. We can find elements $\gamma_p, \gamma_q \in \mathcal{D}$ representing w_p, w_q respectively with $N(\gamma_p) = p$ and $N(\gamma_q) = q$ such that $\gamma_p\tilde{s} = \bar{\tilde{s}}$ and $\gamma_q\tilde{s} = \tilde{s}$. Therefore some powers of γ_p and of γ_q are in $\mathbf{Q}^\times$. Since γ_p and γ_q are quadratic over

$\mathbf{Q}$, it follows $\gamma_p^2 \in \mathbf{Q}^\times$ and $\gamma_q^2 \in \mathbf{Q}^\times$ (see the analysis in [Kur, Proposition 4-4]). Hence $\gamma_p^2 = -p$ and $\gamma_q^2 = -q$. Since γ_q and $\gamma_p\gamma_q\gamma_p^{-1}$ both fix $\tilde{s}$, it follows that $\gamma_p\gamma_q\gamma_p^{-1} = -\gamma_q$ and hence $B^{p\text{-int}} \cong B(-p, -q)$. Conversely suppose $B^{p\text{-int}} \cong B(-p, -q)$. Pick γ_p, γ_q satisfying $\gamma_p^2 = -p$, $\gamma_q^2 = -q$, and $\gamma_p\gamma_q = -\gamma_q\gamma_p$. Let $\tilde{s}$ be the unique edge of Δ reversed by γ_p . Then $\gamma_q(\tilde{s})$ must be $\tilde{s}$, so the image r of $\tilde{s}$ in $G^{(q)}$ is such that $\ell(r)$ is even and $w_p(r) = \bar{r}$. This completes the proof of statement (3) of Theorem 2.

2. PROOF OF THEOREM 1

We now prove Theorem 1. With B, p , and q satisfying the hypotheses of the theorem, let $g(V_B)$ be the genus of V_B . Then by the Riemann-Hurwitz formula the genus g of $V^{(p)}$ is given by

$$(2) \quad g = \frac{g(V_B) + 1}{2} - \frac{e(p)}{4},$$

where $e(p)$ is the number of fixed points of the w_p on $V_B(\mathbf{C})$, cf. [Ogg, Eq. 3]. Now $e(p)$ is equal to the number of points on V_B with complex multiplication by $\mathbf{Z}[\sqrt{-p}]$, which is twice the class number h_p of $\mathbf{Q}(\sqrt{-p})$ if $\mathbf{Q}(\sqrt{-p})$ splits B and 0 otherwise. As q is inert in $\mathbf{Q}(\sqrt{-p})$ we obtain:

$$(3) \quad e(p) = 2h_p \equiv 4 \pmod{8}$$

since $p \equiv 5 \pmod{8}$; see, e.g., [BS, p. 250, Exercise 25]. In particular then $e(p)/4$ is odd. Under our hypotheses Eichler's mass formula gives

$$(4) \quad \frac{g(V_B) + 1}{2} = 1 + \frac{(p-1)(q-1) - 16}{24} \equiv 1 \pmod{2}$$

(see, e.g., [JL2, p. 46]). Combining then (2), (3), and (4), we see that g is even.

From Theorem 2 we have that $\text{Pic}^1 V^{(p)}(\mathbf{Q}_p)^+ \neq \emptyset$ and $\text{Pic}^1 V^{(p)}(\mathbf{R})^+ \neq \emptyset$ since $\mathbf{Q}(\sqrt{p})$ splits B . Furthermore $B^{q\text{-int}} \not\cong B(-1, -pq)$ since $(-1, -pq)_p = +1$ and $B^{q\text{-int}} \not\cong B(-p, -q)$ since $(-p, -q)_q = -1$. Hence $\text{Pic}^1 V^{(p)}(\mathbf{Q}_q)^+ = \emptyset$ by Theorem 2(3). Since $\text{Pic}^2 V_B(\mathbf{Q}_\ell)^+ \neq \emptyset$ for all primes $\ell \leq \infty$ and $\text{Pic}^1 V_B(\mathbf{Q}_\ell)^+ \neq \emptyset$ for primes $\ell \neq p, q, \infty$ by [JL2, Theorem 2], the same holds for $V^{(p)}$. Thus $\text{Pic}^{g-1}(V^{(p)})(\mathbf{Q}_\ell)^+ = \emptyset$ for exactly one prime $\ell \leq \infty$, namely $\ell = q$. By the criterion of Poonen and Stoll, it follows that the jacobian of $V^{(p)}/\mathbf{Q}$ is odd.

It is routine that at most finitely many of the $V_B^{(p)}/\mathbf{Q}$'s in Theorem 1 are hyperelliptic, cf. [Ogg, Sect. 5]. The curves $V_B^{(p)}$ and V_B have good reduction $\tilde{V}_B^{(p)}/\mathbf{F}_2$ and $\tilde{V}_B/\mathbf{F}_2$, respectively, at 2. If $V_B^{(p)}$ were hyperelliptic, we would then have that

$$(5) \quad \#\tilde{V}^{(p)}(\mathbf{F}_4) \leq 2 \cdot \#\mathbb{P}^1(\mathbf{F}_4) = 10.$$

But the supersingular locus on $\tilde{V}_B/\mathbf{F}_2$ is rational over $\mathbf{F}_4$ and its cardinality is the class number h of the definite quaternion algebra of reduced discriminant $2pq$. By Eichler's

formula (see [Kur, (4-1)]) however, we have

$$(6) \quad \#\tilde{V}_B(\mathbf{F}_4) \geq h \geq \frac{(p-1)(q-1)}{12}.$$

Since $\tilde{V}_B$ is a double cover of $\tilde{V}_B^{(p)}$ over $\mathbf{F}_2$, this implies that

$$(7) \quad \#\tilde{V}_B^{(p)}(\mathbf{F}_4) \geq \frac{(p-1)(q-1)}{24}.$$

From (5), (6), and (7) we see that if $V_B^{(p)}$ is hyperelliptic, then $(p-1)(q-1) \leq 240$.

REFERENCES

- [BS] Z. I. Borevich and I. R. Shafarevich, *Number Theory*, Academic Press, New York San Francisco London 1966.
- [Čer] I. V. Čerednik, *Uniformization of algebraic curves by discrete arithmetic subgroups of $\mathrm{PGL}_2(k_w)$ with compact quotients* (in Russian), Math. Sb. (N.S.) **100** (1976) 59–88, Math. USSR Sb. **29** (1976), 55–78.
- [Dr] V. G. Drinfeld, *Coverings of p -adic symmetric regions* (in Russian), Functional. Anal. i Priložen. **10** (1976), 29–40, Functional Anal. Appl. **10** (1976), 107–115.
- [JL1] B. W. Jordan and R. Livné, *Local diophantine properties of Shimura curves*, Math. Ann. **270** (1985), 235–248.
- [JL2] B. W. Jordan and R. Livné, *Divisor classes on Shimura curves rational over local fields*, J. reine angew. Math. **378** (1987), 46 – 52.
- [JLV] B. W. Jordan, R. Livné, and Y. Varshavsky, *Local points on p -adically uniformized Shimura varieties*, in preparation.
- [Kur] A. Kurihara, *On some examples of equations defining Shimura curves and the Mumford uniformization*, J. Fac. Sci. Univ. Tokyo, Sec. 1A **25** (1979), 277–301.
- [Ogg] A. Ogg, *Real Points on Shimura curves*, in: M. Artin and J. Tate, editors, *Arithmetic and Geometry* Volume I, Birkhäuser, Boston Basel Stuttgart, 1983.
- [PSt] B. Poonen and M. Stoll, *The Cassels-Tate pairing on polarized abelian varieties*, preprint, Dec. 1997.

DEPARTMENT OF MATHEMATICS, BOX G-0930, BARUCH COLLEGE, CUNY, 17 LEXINGTON AVENUE, NEW YORK, NY 10010 USA

MATHEMATICS INSTITUTE, THE HEBREW UNIVERSITY OF JERUSALEM, GIVAT RAM, JERUSALEM 91904 ISRAEL

E-mail address: rlivne@sunset.ma.huji.ac.il